

Upper limit of the energy of the photon and the phase change of photon to material particles at the Schwarzschild radius

C. Radhakrishnan Nair
Institute of Mathematical Sciences
Kariyavattom
Thiruvananthapuram
Pin 695 581
India
E-mail: nairrcc@rediffmail.com

Abstract

The concept of Schwarzschild radius is extended to the photon and the upper limit imposed on the energy of a photon as a result of the three characteristics of the photon—the constancy of the velocity of light, the spin value of $1\hbar$ and the zero rest mass of the photon—is shown. Further the phase change that occurs to the photon at the Schwarzschild radius, from energy to matter as a result of vacuum fluctuations is indicated.

The photon, as an elementary particle, is unique. It is the only elementary particle of energy whereas there are hundreds of elementary particles of matter. There are three unique properties for the photon.

1. In the free space, the photon always propagates with the same constant velocity, velocity of light c , irrespective of the energy it possesses.
2. The rest mass of the photon is zero.
3. The photon has a constant spin value of $\hbar$

In this short paper, we shall examine one remarkable consequence of the above three properties of the photon.

The special theory of relativity postulates an upper limit [1] for the velocity of light in free space. We shall show that the upper limit of the velocity of light puts an upper limit on the rotational angular velocity and hence on the total energy possible for a photon.

Relativistic equation for the energy of a particle is given by the equation

$$E^2 = p^2 c^2 + m_0 c^2 \quad (1)$$

where

E = Energy of the particle

p = Momentum of the particle

c = velocity of light

m_0 = Rest mass of the particle.

For a photon, the rest mass is zero and therefore the energy of the photon is given by the equation

$$E^2 = p^2 c^2 \quad (2)$$

It is well known that photon has one degree of freedom of rotation and two degrees of freedom of translation [2, 3]. Since the photon has no rest mass, the total energy of the photon must be stored as rotational kinetic energy and translational kinetic energy.

The first postulate of relativity [1] (or the Galilean invariance) states that no object is aware of its motion when moving with uniform velocity and light is no exception to this. Therefore, for the photon itself, moving with the velocity of light, its kinetic energy due to translatory motion will be zero. But this does not say that the photon does not exist for itself. The photon has a kinetic energy of rotation and therefore the photon can feel its existence. Incidentally, this also shows the necessity of a nonzero spin value for any particle with zero rest mass. Therefore, from the point of view of the photon itself, since the photon has no rest mass, the total energy of the photon is stored as rotational kinetic energy. We can associate rotation even to a particle with zero rest mass, provided the particle has spatial extension.

The energy of the photon is given by the expression

$$E = h\nu = \frac{hc}{\lambda} \quad (3)$$

where h = the Planck's constant, ν = the frequency of the photon, λ = the wavelength of the photon.

The wavelength guarantees the extension of the photon. The expression $E = \frac{hc}{\lambda}$ shows that the greater the energy of the photon, the shorter its wavelength. This implies that with the increase in the energy, the photon will be more and more localized. Can this localization go on ad infinitum? The idea of infinite amount of energy packed in an infinitely small volume is quite unphysical and it cannot be true. How is the property of the space altered by the ever increasing energy density? Do the constant parameters $\hbar$ and c put an upper limit for the possible energy of a photon? What are the consequences of this upper limit on the space time geometry?

The Harward Tower experiment [4] has clearly demonstrated that the energy of a photon is increased under gravitational attraction. Can the energy of the photon be increased continuously without limit by gravitational attraction? Since the photon itself cannot feel any increase in the translational kinetic energy, the increased energy of the photon, when attracted by a gravitational field, is entirely stored as rotational kinetic energy. Spin is the quantum mechanical analogue of classical angular momentum. Since the photon has a spin of $1 \hbar$, we can attribute an angular momentum to the photon in the semiclassical analysis.

When L , I and w are the angular momentum, moment of inertia and angular velocity of a particle, we have the expression

$$L = Iw \quad (4)$$

When m , r and v are the mass, radius and velocity of the particle respectively, we have

$$L = mrv \quad (5)$$

Special theory of relativity puts c as the upper limit for the velocity of any signal. Therefore no particle can rotate with a tangential linear velocity v greater than the velocity of light. When we put this limiting value of c to v in equation (5), we have

$$L = mrc$$

When the particle we consider is the photon, L is replaced by the spin S of the photon.

$$S = mrc \quad (6)$$

For a photon $S = 1 \hbar$

$$\therefore 1 \hbar = mrc.$$

Here m is the equivalent mass of the photon due to its spin.

$$m = \frac{\hbar}{rc} \quad (7)$$

We assume that as in the case of material particles, the concept of Schwarzschild radius is valid for the photon also.

The Schwarzschild radius of a black hole is given by the expression

$$r = \frac{2Gm}{c^2} \quad (8)$$

$$\frac{r}{m} = \frac{2G}{c^2} \quad (9)$$

The right hand side of the equation (9) is a constant. Therefore, for any black hole, the increase in the radius must be accompanied by an increase in the mass. But when we consider the photon as a black hole, Schwarzschild radius is the minimum radius that a photon can have. In the case of material particles, we can have black holes of varying masses with the corresponding radii. This is possible because the right hand side value of equation (9) can be maintained by proportional increase in the values of r and m . This is impossible for the photon because for the photon, with the increase in the equivalent mass, the radius decreases. Therefore, unlike in the case of material particles, there is a unique value of radius and mass, for a photon to behave as a black hole.

Substituting the value of the radius r from equation (8) into equation (7), we get

$$m = \sqrt{\frac{\hbar c}{2G}} \quad (10)$$

This is the expression for Planck mass. Planck mass is usually defined as the minimum mass corresponding to the Schwarzschild radius of a classical particle. Planck mass, in the case of the photon, is the maximum equivalent mass permitted for the photon.

We have

$$\begin{aligned} c &= 2.998 \times 10^8 \text{ m s}^{-1} \\ h &= 6.62 \times 10^{-34} \text{ Js} \\ G &= 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2} \end{aligned}$$

Substituting these values we get

$$m = 1.54 \times 10^{-7} \text{ kgs}$$

The maximum energy possible for the photon will be

$$\begin{aligned} mc^2 &= \sqrt{\frac{\hbar c}{2G}} c^2 \\ &= 8.61 \times 10^{22} \text{ MeV.} \end{aligned} \tag{11}$$

The highest energy so far observed [5] for a photon is of the order of 10^{13} eV.

A simple dimensional analysis in terms of the Planck's natural units will be illuminating here. The dimension of energy is ML^2T^{-2} . When we substitute for M , L and T in terms of c , G and h we get the expression

$$\begin{aligned} E &= ML^2T^{-2} \\ &= \sqrt{\frac{\hbar c}{G}} \frac{\hbar G}{c^3} \frac{c^5}{\hbar G} \\ &= \sqrt{\frac{\hbar c}{c^2}} \end{aligned} \tag{12}$$

Expressions (11) and (12) are identical except for the geometrical factor of 4π appearing in equation (11).

What happens to the photon when the photon reaches its upper limit of energy, having an equivalent mass equal to the Planck mass. Energy and matter are two phases of the same physical entity. Hawking [6, 7] has shown that in the case of primordial black holes, at the Schwarzschild radius, spontaneous pair production takes place. At the upper limit of energy permitted for the photon, having reached the Schwarzschild radius, it may be legitimate to presume that the photon will undergo pair production and transformed to material particles. It is also an open question whether the photon at the maximum energy limit can be disintegrated into many photons with smaller energies and larger wavelengths in the turbulent sea of vacuum fluctuations. Of course these are pure speculations, either to be proved or disproved by future theoretical and experimental studies.

Acknowledgement

Discussions with G. Mohanachandran, T. E. Gireesh, K. Suresh Babu and M. Sabir were extremely useful. L. A. Ajith has happily helped me in the

preparation of the script.

References

- [1] Einstein A. (1905) *Ann. Phys. Lpz.* **17**, 140.
- [2] Han D., Kim Y. S. and Son D. (1982) *Phys. Rev. D* **25**, 461.
- [3] Kim Y. S. and Wigner E. P. (1990) *J. Math. Phys.* **31**, 55.
- [4] Pound R. V. and Rebka, Jr G. A. (1960) *Phys. Rev. Lett.* **4**, 337.
- [5] Horan D. and Weekes T. C. (2004) *New. Astro. Rev.* **48**, 527.
- [6] Hawking S. W. (1974) *Nature* **248**, 30.
- [7] Hawking S. W. (1975) *Commun. Math. Phys.* **43**, 199.